

Theory and observation of frequency splitting and sweeping in tokamaks

R. G. L. Vann¹, R. O. Dendy^{2,1}, M. P. Gryaznevich² and S. E. Sharapov²

¹*Department of Physics, University of Warwick, Coventry CV4 7AL, U.K.*

²*Euratom/UKAEA Fusion Association, Culham Science Centre, Abingdon, Oxfordshire OX14 3DB, U.K.*

Frequency splitting and frequency sweeping (“chirping”) of high frequency MHD modes are widely observed in tokamak plasmas [1,2]. Here we consider observations of frequency splitting on JET (see for example Fig. 1) and chirping on MAST (Fig. 3), and show how these may be modelled by the Berk-Breizman augmentation of the Vlasov-Maxwell equations (henceforth “the VM(BB) system”) [3–5]. The VM(BB) system models the coupling between energetic particles and the wave modes they excite, based on the one-dimensional electrostatic bump-on-tail problem with particle distribution relaxation and background electric field damping. We cast the model as the follows [7], in terms of the particle distribution $f(x, v, t)$ and the electric field $E(x, t)$:

$$\frac{\partial f}{\partial t} + v \frac{\partial f}{\partial x} + E \frac{\partial f}{\partial v} = -\nu_a (f - F_0) \quad (1)$$

$$\frac{\partial E}{\partial t} + \int v (f - f_0) dv = -\gamma_d E \quad (2)$$

Here F_0 denotes the combined particle source and loss function, ν_a the particle relaxation rate, γ_d the combined effect of all background damping mechanisms that act on the electric field, and f_0 the spatial mean of f . Spatial lengths are normalised to the Debye length λ_D ; velocities to the thermal speed v_{the} ; time to the inverse plasma frequency $\omega_p^{-1} \equiv \lambda_D/v_{the}$; and E to $m_e v_{the}^2/e\lambda_D$.

A code has recently been developed [6] that allows direct numerical solutions of the fully nonlinear VM(BB) system across the entirety of (γ_d, ν_a) parameter space for any $F_0(v)$. Application of this code [7] for a particular $F_0(v)$ shows how the behaviour of the VM(BB) system depends on its parameters. We can take a cut through (γ_d, ν_a) parameter space and observe how the behaviour of the system varies as we travel along this line. Consider the line $\gamma_d = 1.0$: for each value of ν_a , we extract the extreme values achieved by the electric field energy (excluding the initial transient phase), and plot in Fig. 2 these extrema as a function of ν_a . For example, sinusoidal behaviour at a particular ν_a would contribute two points to the plot. Provided the underlying period varies slowly with ν_a , an abrupt splitting of an observed extremum into two branches implies a period doubling bifurcation at that parameter value. Figure 2 demonstrates how the VM(BB) system naturally generates frequency splitting phenomenology as seen experimentally in Fig. 1.

Figure 1: Experimental observation of frequency splitting. This plot shows TAEs in JET shot 40332 undergoing period doubling bifurcations (reported and discussed in [4]).





Figure 2: Plots of extrema in the electric field energy $A(t)$ observed in the solution of the VM(BB) system as a function of particle relaxation rate ν_a (the parameter γ_d is fixed at unity). The splitting of extrema corresponds to period doubling bifurcations in the time series $A(t)$. The plot on the right is a magnification of a section of the plot on the left; it shows the bifurcation path of the system to chaos through a series of period doublings.

Figure 3 shows MAST data obtained from a directional decomposition of 1MHz signals observed at three midplane Mirnov coils each separated by a toroidal angle of 60° . The high frequency MHD mode shown in Fig. 3 is identified as a TAE excited by the energetic ion population and reacting back upon it [8]. We now show how this class of self-consistent evolution can be modelled by numerical solution of the fully nonlinear VM(BB) system.

Figure 4 shows the time evolution of the frequency of the first spatial mode for the fully nonlinear BB system. This plot is obtained by taking fast Fourier transforms on successive Hanning windows. Here the two control parameters, ν_a and γ_d , are fixed throughout the simulation, corresponding to a quasistationary background plasma and energetic particle drive. Time evolution of the system thus arises solely from the self-consistent interaction of the energetic particle population and its associated field. Like

Fig. 3, Fig. 4 exhibits repeated bursts whose frequency undergoes almost-symmetric up-down chirping. Direct measurements of the spatially averaged distribution function $f_0(v)$, linked to corresponding instants during the evolution of a single burst, are shown in Fig. 5, which demonstrates the central role of hole-clump pair formation and evolution. This is entirely responsible for the chirping seen in Fig. 4, which suggests that this may be the mechanism underlying the phenomenology observed in the real tokamak plasma (Fig. 3).




Figure 3: Experimental observation of frequency chirping in nine successive bursts of TAE activity. Magnitude of MHD activity measured in neutral beam-heated MAST pulse 5568 during a 13ms interval, showing frequency in the range 80-120kHz.




Figure 4: Frequency chirping in successive bursts of activity from the fully nonlinear VM(BB) model. Dimensionless time and frequency are normalised by ω_p . The plot shows mode amplitude on a logarithmic colour scale.

Our novel splitting and sweeping results provide fresh evidence of the range of tokamak phenomenology that is captured by the fully nonlinear VM(BB) model.




Figure 5: Hole-clump pair formation and evolution shown at four different times during the simulation whose spectrogram is shown in Fig. 4. Available as a movie [9].

We thank T. Arber, B. Breizman and H. Berk for helpful discussions. This work was supported in part by the United Kingdom Engineering and Physical Sciences Research Council and by Euratom. Computing facilities were provided by the Centre for Scientific Computing at the University of Warwick with support from Joint Research Equipment Initiative grant JR00WASTEQ.

References

- [1] A. Fasoli *et al.*, *Plasma Phys. Control. Fusion* **39**, B287 (1997)
- [2] M. P. Gryaznevich, *presented at 8th IAEA TCM on EPs* (2003) <http://fusion.gat.com/conferences/iaea-tm-energetic/>
- [3] H. L. Berk, B. N. Breizman, and M. Pekker, *Phys. Rev. Lett.* **76**, 1256 (1996)
- [4] A. Fasoli *et al.*, *Phys. Rev. Lett.* **81**, 5564 (1998)
- [5] H. L. Berk, B. N. Breizman, J. Candy, M. Pekker, and N. V. Petviashvili, *Phys. Plasmas* **6**, 3102 (1999)
- [6] T. D. Arber and R. G. L. Vann, *J. Comput. Phys.* **180**, 339 (2002)
- [7] R. G. L. Vann, R. O. Dendy, G. Rowlands, T. D. Arber, and N. d'Ambrumenil, *Phys. Plasmas* **10**, 623 (2003)
- [8] H. L. Berk *et al.*, *Proc. 31st EPS Conf. Plasma Phys.* (2004)
- [9] http://www.warwick.ac.uk/staff/Roderick.Vann/research/hole_clump.html