

ABELISATION OF THE NEUTRON PROFILE DATA AT JET USING MINIMUM FISHER REGULARISATION

J. Mlynar¹, G. Bonheure², A. Murari³, S. Popovichev⁴, V.Svoboda⁵

and JET EFDA contributors*

¹Association Euratom-IPP.CR, Za Slovankou 3, CZ-182 00 Prague 8, Czech Republic

²Association Euratom-Belgian state, Partners in the TEC, ERM-KMS, Brussels, Belgium

³Consorzio RFX-Associazione EURATOM-ENEA per la Fusione, I-35127 Padova, Italy

⁴Euratom/UKAEA Fusion Association, Culham Science Centre, Abingdon OX14 2DB, UK

⁵Association Euratom-IPP.CR, FJFI CVUT, Brehova 7, CZ-115 19 Prague 1, Czech Republic

*See the Appendix of M.L.Watkins et al., Fusion Energy 2006, Proc. 21st Int. Conf. Chengdu

INTRODUCTION

Analyses and validation of data from neutron measurements present one of the key contributions of JET to research and development of diagnostic systems for fusion burning plasmas. In this respect the JET neutron profile monitor has a unique importance in determining spatial distribution of neutron emissivities [1]. The monitor consists of two cameras, one with ten collimated channels, and another with nine collimated channels providing fan-shaped horizontal and vertical views, respectively. The spatial plasma coverage provided by the neutron profile monitor is adequate for inverse reconstructions, albeit with a rather scarce spatial resolution. Minimum Fisher Regularisation (MFR) has been recently applied at JET for the 2D tomographic reconstruction of the neutron emissivity [2]. This contribution presents the latest extension of the algorithm that allows for emissivity profile reconstruction (abelisation) under the assumption of constant neutron emissivities along magnetic flux surfaces (i.e. full poloidal symmetry). This strong assumption, when realistic, allows for profile analyses even if neutron emissivities are low. The MFR abelisation principle is tested and validated on phantom functions, including simulations of emissivity evolution and random data noise. First analyses of experimental data are presented, with particular interest in their potential use for tritium transport studies.

ABELISATION VIA RAPID MINIMUM FISHER REGULARISATION

Plasma tomography can retrieve emissivities g_j from line integrals f_i :
$$f_i = \sum_j^N T_{ij} g_j + \zeta_i$$

where T_{ij} is geometric matrix, N number of pixels, ζ_i existing statistical and systematic errors. In abelisation, g_j is supposed to be axisymmetric, i.e. pixels are delimited by magnetic flux surfaces. In the MFR abelisation the matrix T_{ij} corresponds to the total length of the i -th viewing line in the j -th pixel. A smooth solution for g_j is found by minimising

$$\Lambda_{MF} = \frac{1}{2} \chi^2 + \alpha_R I_F, \text{ where } \chi^2 = \frac{1}{M} \sum_i \left(\frac{\varepsilon_i}{\sigma_i} \right)^2 ; \varepsilon_i = f_i - \sum_j T_{ij} g_j \text{ is the "retrofit"; } \alpha_R \text{ is a}$$

regularisation (smoothing) parameter, σ_i the data errorbars, M number of viewing lines and I_F the Fisher Information of the emissivity distribution. Minimal I_F corresponds to emissivity distribution estimate with minimum variance. In practice this principle leads to higher smoothness in low emissivity regions. These postulates have been implemented into a Tikhonov regularisation algorithm, see [3]. A rapid version can simultaneously proceed data with their time characteristics [4] providing an efficient tool for studies of the emissivity evolution. Spatially and temporarily resolved neutron emissivities can be subsequently decomposed into spatial eigen-modes (topos) and corresponding time eigen-vectors (chronos) using Singular Value Decomposition (SVD): $g(r, t) = \sum_i v_i(r) s_i u_i(t)$, for details see [3].

TESTS OF THE MFR ABELISATION ON PHANTOM EMISSIVITIES

The rapid MFR Abelisation algorithm was tested and optimised in runs with simple phantom functions that model diffusion-like evolutions of neutron emissivities, see e.g. Fig. 1a. In order to keep the tests realistic, the geometric matrix corresponded to experimentally determined flux coordinates (JET pulse #61161, $t = 10.5$ s) and $> 3\%$ random noise was




Figure 1: Tests of the MFR abelisation on projections of phantom data with 3% added random noise
a) original phantom function b) example of the reconstructed function c) time averaged relative differences (reconstruction errors) for 20 different runs of the MFR abelisation

superposed onto the phantom projections. Within the tests the MFR abelisation was optimised and validated however with a conclusion that central regions tend to be either oversmoothed or noisy, see Fig. 1b. Low spatial resolution was identified as the main cause: Only two vertical and two horizontal chords carry information on emissivities from the $r/a < 0.2$ region. Due to this feature it currently seems difficult to fully automate the MFR abelisation. Errors of the inverse reconstruction can be determined via simplified Monte Carlo simulations: Results are presented in Fig. 1c, where dots correspond to time averaged errors in 20 consecutive reconstructions of the phantom projections that differed only in the superposed 3% random noise. Other tests proved that the SVD analyses lead to highly dependent results even with noisy data.

EXPERIMENTAL RESULTS

In [5] it has been shown that at JET, tritium puff into a low density plasmas leads to D-T (14 MeV) neutron emissivities with good poloidal symmetry. The corresponding data are therefore suitable for abelisation, which is the case e.g. in JET pulse #61161. Rapid MFR abelisation of this event, presented in Fig. 2a, offered best results for α_R corresponding to 4% expected errors in central chords. Evolution of the reconstructed D-T neutron emissivity profile clearly visualise the Tritium influx, which is even more apparent in the consecutive SVD of the reconstruction, see Fig. 2b. These results can be used as input for the ratio method [5] which can lead to quantitative estimates of transport parameters. Interestingly, retrofit of the reconstruction on the input data unveils a minor assymetry, with a slightly higher signal




Figure 2: Experimental results – studies of Tritium puff: a) reconstructed evolution of emissivity profiles of 14 MeV neutrons b) first three orders of the Singular Value Decomposition of the reconstructed emissivity evolution

towards the high field side (!). Notice that the assymetries may originate in offsets of fast particles and/or in minor errors of the magnetic reconstruction [4]. The MFR abelisation was also applied on data from recent JET campaign that focussed on magnetic ripple studies: D-D (2.5 MeV) neutron emissivity in JET pulse #69604 (low ripple) was compared to emissivity in a quasi-identical pulse #69610 (high ripple) at $t = 21.5$ s. Due to low neutron intensities in these pulses, neutron data for abelisation had to be integrated over one second. Nonetheless, in order to get a smooth reconstruction the factor α_R had to be high ($\sigma \sim 8\%$ in central chords). The retrofit identified that this condition was caused by apparent data assymetry in the vertical channels towards low-field side. Still, the results of MFR abelisation clearly demonstrate - as expected - that the neutron emissivity profile is broader in plasmas with higher ripple.

To conclude, the MFR abelisation of neutron emissivities at JET proved to be feasible, albeit with limitations in the central plasma region. However, most of the data show some degree of poloidal assymetry, so that the MFR abelisation will be probably limited to a secondary role in the ongoing neutron data analyses. Importantly, the MFR abelisation can provide a precise indication on poloidal data asymmetries. The results of this study give strong incentive for further development of the 2D tomography.

ACKNOWLEDGEMENTS

This work was carried out under the European Fusion Development Agreement.

REFERENCES

- [1] A. Murari et al, "Burning plasma" diagnostics for the physics of JET and ITER, Plasma Phys. Control. Fusion 47 (2005) B249
- [2] G. Bonheure et al, 2-D Spatial Distribution of D-D and D-T Neutron Emission in JET ELMY H-mode Plasmas with Tritium Puff, Proc. 32nd EPS Conference Tarragona, P1-083 (2005)
- [3] M. Anton et al., X-ray tomography on the TCV tokamak, Plasma Phys. Control. Fusion 38 (1996) 1849-1878
- [4] J. Mlynar et al, Investigation of the consistency of magnetic and soft X-ray plasma position measurements on TCV by means of a rapid tomographic inversion algorithm, Plasma Phys. Control. Fusion 45(2003) 169-180
- [5] G. Bonheure et al., Neutron profiles and fuel ratio n_T/n_D measurements in JET ELMY H-mode plasmas with tritium puff, Nucl. Fusion 46 (2006) 725