

Nonlinear gyrokinetic simulations of microtearing mode turbulence

W. Guttenfelder¹, J. Candy², S.M. Kaye¹, W.M. Nevins³, E. Wang³, R.E. Bell¹,
G.W. Hammett¹, B.P. LeBlanc¹, D.R. Mikkelsen¹, H. Yuh⁴

¹Princeton Plasma Physics Laboratory, Princeton NJ 08543 USA

²General Atomics, San Diego, CA 92186 USA

³Lawrence Livermore National Laboratory, CA 94551 USA

⁴Nova Photonics Inc., Princeton, NJ 08540 USA

First non-linear gyrokinetic simulations of microtearing mode turbulence have recently been reported for a high-collisionality NSTX discharge ($B_T=3.5\text{ kG}$, $I_p=0.7\text{ MA}$, $R/a=0.82/0.62\text{ m}$) [1] and independently for conventional tokamak parameters [2]. Using the Eulerian gyrokinetic code GYRO [3] with finite collisionality ($v_{ei}=1.26\text{ c}_s/a$, $Z_{eff}=2.9$) and electromagnetic perturbations ($\delta A_{||}$, $\beta_e=8\pi n_e T_e / B_0^2 = 8.8\%$), it was shown for NSTX at $r/a=0.6$ (where only microtearing modes are linearly unstable) that relatively fine radial grid spacing was required to sufficiently resolve fluctuations and demonstrate monotonically decaying turbulence spectra [1]. Fig. 1 shows a snapshot of the resulting density ($\delta n/n_0$) and magnetic field (δB_r) fluctuations at a fixed toroidal angle in real space (R, Z) coordinates. Narrow




Fig. 1. (left) Normalized density perturbations and (right) shear magnetic field perturbations.

radial density perturbations are apparent on the outboard side, which correspond to the narrow resonant parallel current perturbations responsible for destabilizing the microtearing mode. To resolve these perturbations, the simulation used 400 radial grid points [and 8 complex toroidal modes with toroidal mode spacing $\Delta n=5$ or $(k_0\rho_s)_{\min}=0.105$, $k_0=nq/r$, $q=1.69$], with a spacing $\Delta x=0.2 \rho_s$ that is capable of resolving the high order rational surfaces ($n_{\max}=35$, $k_0\rho_{s,\max}=0.735$) separated by $\Delta r_{\text{rat},\min}=1/n_{\max}q'=1/k_{0,\max}s=0.78\rho_s$ ($s=r/q \cdot dq/dr=1.75$). An additional simulation using 16 toroidal modes and 540 radial grid points ($\Delta n=3$, $\Delta x=0.15 \rho_s$), requiring $\sim 250,000$ cpu-hrs on the Oak Ridge “Jaguar” Cray XT5, exhibits very similar behavior, illustrating, at least qualitatively, the microtearing turbulence is resolved.

While local peak density perturbations of $\delta n/n \sim 2\%$ occur around the entire poloidal circumference, the δB_r perturbations are strongly ballooning. The structure of δB_r is spatially broad on the outboard midplane with local instantaneous values as large as 30 Gauss ($\sim 1\%$ of the vacuum field strength, $B_0=3.5\text{kG}$). Using both δn and δB_r from multiple time slices, calculations predict $1-2^\circ$ of instantaneous polarimetry phase shift ($\sim 0.3^\circ$ rms) dominated by δB_r , which may be measurable by a new polarimetry system to be installed on NSTX [4]. Furthermore, the outboard density perturbations may be measurable by the “high-k” coherent scattering diagnostic [5] which is sensitive to density perturbations with $k_r > k_\theta$.

The magnetic “flutter” from the strong δB_r fluctuations is responsible for $\sim 98\%$ of the total electron thermal transport, $\chi_{e,\text{sim}} \approx 1.2 \rho_s c_s^2/a = 6 \text{ m}^2/\text{s}$, which is in the experimental range $\chi_{e,\text{exp}} = 5-8 \text{ m}^2/\text{s}$. Estimates using the spectrum of rms saturated amplitudes ($\langle |\delta B_r/B|^2 \rangle^{1/2} \sim 0.15\%$) predict that island widths, $w_{\text{island}} = 4\sqrt{\delta B_r/B \cdot rR/\text{ns}}$, should be larger than the minimum separation in rational surfaces $w_{\text{island}}/\delta r_{\text{rat}} \leq 8$, $\delta r_{\text{rat}} \approx \Delta n/n^2 q'$ [1], satisfying the Chirikov overlap criteria [6]. Poincare surface-of-section plots, formed by integrating 100 field line trajectories for 3000 poloidal transits, confirm global stochasticity throughout the entire simulation domain [7]. From the integrated trajectories the magnetic diffusivity is calculated $D_M \approx 6 \times 10^{-7} \text{ m}$, from which we infer a magnetic correlation length $L_c \approx 0.25 \text{ m}$ much shorter than the electron mean free path $\lambda_{\text{mfp}} = 12.5 \text{ m}$. The resulting collisionless test particle stochastic transport model, $\chi_{e,\text{st}} = 2\sqrt{2/\pi} \cdot f_p D_m v_{Te}$, [7] agrees to within $\sim 20\%$ of the simulated transport.

Fig. 2 shows the predicted transport from simulations that vary normalized electron temperature gradient (a/L_{Te}), collisionality, beta, and $E \times B$ shear (γ_E , defined in [8]) using the smaller of the above grids ($\sim 100,000$ cpu-hrs/sim). The electron thermal transport is very “stiff”, varying $\pm \sim 100\%$ for $\pm 20\%$ changes in a/L_{Te} , suggesting it would be useful to characterize the scaling of the threshold gradient. Most noteworthy is that the predicted collisionality scaling, $\chi_e \sim v_e^{1.1}$, is roughly consistent with global energy confinement scaling in both NSTX and MAST [9], $\Omega_i \tau_E \sim v_*^{-(0.82-0.95)}$. This gives strong evidence for the importance of microtearing modes in determining confinement scaling in NSTX, as no other micro-instability is predicted to be unstable for these parameters. However, including the experimental value of $E \times B$ shear, $\gamma_{E,exp} \approx 0.17$ c_s/a (comparable to the maximum linear growth rate) reduces the predicted transport to negligible values. Given the stiff behavior with a/L_{Te} , an additional γ_E scan was run with $1.2 \times a/L_{Te,exp}$. The transport at $\gamma_{E,exp}$ is now larger, but still




Fig. 2. Predicted χ_e vs. (a) a/L_{Te} , (b) $v_{ei} (c_s/a)$ (log-log scale), (c) β_e , and (d) $E \times B$ shear rate, $\gamma_E (c_s/a)$. The shaded regions indicate experimental values with uncertainties.

3-4× smaller than experiment. While the radial resolution in these simulations is marginally sufficient to distinguish the narrow resonant parallel current perturbations ($\Delta_j \leq 0.3 \rho_s$) responsible for the microtearing instability, the quantitative linear growth rates are smaller than highly resolved linear simulations ($\Delta x \approx 0.03 \rho_s$ for $k_0 \rho_{s,\max}$), especially as marginal stability is approached (e.g. reduced a/L_{Te} , v_e , or β_e). As a result, with improved non-linear resolution the magnitude of transport in Fig. 2 might be increased sufficiently to recover experimental transport with $\gamma_{E,exp}$. Another uncertainty in predicting *quantitative* transport stems from the limited resolution in toroidal modes (k_0). Fig. 3 shows the fractional transport spectra for three values of γ_E . While there is a well defined peak at $k_0 \rho_s \approx 0.21$ with no $E \times B$




Fig. 3. Fractional transport spectra for three different $E \times B$ shear rates.

with no variation in equilibrium profiles (q , a/L_{Te} , etc...). Including profile variation, which is likely important at large $\rho_s/a \approx 1/120$ in NSTX, may produce some non-local modification of these results.

This work was supported by DOE contracts DE-AC02-09CH11466, DE-FG03-95ER54309, and DE-AC52-07NA27344. We acknowledge generous allocations at NERSC, and at the Oak Ridge Leadership Computing Facility (DOE contract DE-AC05-00OR22725).

References

- [1] W. Guttenfelder, J. Candy, S.M. Kaye, W.M. Nevins, E. Wang et al., Phys. Rev. Lett. **106**, 155004 (2011).
- [2] H. Doerk, F. Jenko, M.J. Pueschel and D.R. Hatch, Phys. Rev. Lett. **106**, 155003 (2011).
- [3] J. Candy & R.E. Waltz, Phys. Rev. Lett. **91**, 045001 (2003).
- [4] J. Zhang et al., Rev. Sci. Instrum. **81**, 10D519 (2010); APS-DPP, Chicago (2010).
- [5] D.R. Smith et al., Rev. Sci. Instrum. **79**, 123501 (2008).
- [6] G.M. Zaslavsky and B.V. Chirikov, Sov. Phys. Usp. 14, 549 (1972).
- [7] E. Wang et al., Phys. Plasmas **18**, 056111 (2011); W.M. Nevins et al., Phys. Rev. Lett. **106**, 065003 (2011).
- [8] R.E. Waltz and R.L. Miller, Phys. Plasmas **6**, 4265 (1999).
- [9] S.M. Kaye et al., Nucl. Fusion **47**, 499 (2007); M. Valovič et al., IAEA EXC/P8-18, Daejon (2010).