

## Rotation-induced electrostatic-potential and its effect on the radial electric field, impurity density and radiated power asymmetries in NSTX

L. F. Delgado-Aparicio<sup>1</sup>, R. E. Bell<sup>1</sup>, G. J. Kramer<sup>1</sup>, M. Podestà<sup>1</sup>, B. P. LeBlanc<sup>1</sup>, A. Diallo<sup>1</sup>, B. Stratton<sup>1</sup>, S. Paul<sup>1+</sup>, S. Kaye<sup>1</sup>, S. Gerhardt<sup>1</sup>, J. Menard<sup>1</sup>, H. Yamazaki<sup>1</sup>, Y. Takase<sup>2</sup> and M. Ono<sup>1</sup>

<sup>1</sup>Princeton Plasma Physics Laboratory, Princeton, NJ, 08540, USA

<sup>2</sup>The University of Tokyo, Graduate School of Frontier Sciences, Kashiwa 277-8561, Japan

### I. Introduction and motivation.-

With the selection of molybdenum and tungsten for the divertor in fusion experiments, the importance of understanding the sources, transport and confinement of high-Z impurities has become crucial. In particular, controlling Z-transport to avoid core accumulation is necessary to achieve and maintain high fusion performance. Along these lines, understanding poloidal asymmetries and its role for reducing the impurity peaking is highly desirable for the development of the spherical tokamak (ST) concept. An important impurity transport task for NSTX-U will be to characterize the properties of H-mode scenarios with high toroidal velocities, in which the ‘standard’ neoclassical theory [1] derived for ‘subsonic’ Mach numbers may no longer be applicable [2-4]. The main motivation behind the installation of the new tangential off-axis neutral-beam injectors in NSTX-U [see Fig 1-a)] was to increase the available heating power, the NBI-driven current and the bootstrap fraction. However, the use of tangential off-axis beams may bring a concomitant increase of the deuterium and impurity Mach numbers providing additional momentum to the outward low temperature plasma [see Fig 1-b)]. In this scenario, the formation of a rotation-induced electrostatic potential ( $\Delta\varphi$ ) with changes in the radial electric field ( $E_r$ ) and the appearance poloidal density asymmetries [2-4] are nearly assured. The computation of these electrostatic potentials is currently being used to study the associated two-dimensional distribution of impurity density and radiated power asymmetries in NSTX and NSTX-U. The Mach number correction enhancing the impurity asymmetries is proportional to the ion-mass so the effects on high-Z impurities such as Fe<sub>26</sub>, Mo<sub>42</sub> and/or W<sub>74</sub> will be easily distinguishable from that of Li<sub>3</sub>, Be<sub>4</sub>, B<sub>5</sub> or C<sub>6</sub>. It has also been theorized that formation of these potential differences and subtle changes in  $E_r$  can possibly vary the heat and particle transport by increasing the fraction of trapped particles [3] and modifying the linear/non-linear stability of gradient-driven modes due to changes in the  $\mathbf{E}\Delta\varphi\times\mathbf{B}$  shearing rates [5], but these findings remain to be evaluated experimentally.

### II. Physics.-

The main centrifugal effect of toroidal rotation on heavy impurities is their radial displacement to the outer plasma, which result in a non-uniform distribution around a magnetic




Fig. 1. NSTX-U’s tangential NBI system and expected torque.




Fig. 2.  $P_{\text{rad}}$  shortfall and in/out asymmetry in NSTX.




Fig. 3. Mid-plane MPTS and CHERS data used as inputs in the calculation.

on electrons, main ions and impurities, a differential electrostatic potential is generated to satisfy quasi-neutrality [3,4]. The calculations presented in this paper start with the mid-plane data measured by the Multi-point Thomson Scattering (MPTS) and Charge Exchange Recombination Spectroscopy (CHERS) diagnostics (see Fig. 3) but relies on the appropriate mapping of flux-surface quantities like electron and ion temperature ( $T_{e,c}$ ) and rotation frequency ( $\omega_\phi$ ) to find the 2D electron, deuterium and carbon density profiles self-consistently assuming the presence of a poloidal variation due to centrifugal forces. The few assumptions considered include a zero electron mass, a D-D plasma, fully-stripped carbon ions, a trace impurity with charge “Z” given by coronal equilibrium ( $\langle Z \rangle(T_e)$ ) and equilibrated ion temperatures (e.g.  $T_D = T_C = T_Z$ ). As result of this numerical procedure nested electron and ion temperatures are mapped on flux surfaces  $\psi(R,Z)$  while the electron density ( $n_e$ ) shows a small departure from poloidal symmetry due to centrifugal effects (see Figure 4).

### III. Electrostatic potential and changes of the radial electric field ( $E_r$ ).

The ideal, *ah-hoc* and numerical iterative solutions for the electrostatic potential difference [ $\Delta\varphi = \varphi(\theta) - \varphi(\theta=0)$ ] are now routinely obtained as depicted in Fig 5. The ideal electrostatic potential shown in Fig. 5-a) is used only as a reference since it is obtained assuming that the main intrinsic impurity (e.g. carbon) is at the trace limit ( $\alpha_C = 36n_C/n_e \ll 1$ ),

$$\Delta\varphi|_{\alpha_C \ll 1} = \frac{T_e}{T_e + T_i} \frac{m_i \omega_\phi^2}{2e} (R^2 - R_0^{*2}) \quad (1)$$

where,  $\omega_\phi$  is the toroidal rotation frequency profile and  $R_0^*$  are the reference positions at the LFS mid-plane. In this limit, the minimum of the  $\Delta\varphi(\theta=\pi)$  is approximately -70 V. The *ad-hoc* solution [5] shown in eq. (2) attempts to extend the previous ideal approximation beyond the carbon trace limit [2] by assuming a finite effective plasma mass ( $m_{eff}$ ) and an effective charge ( $Z_{eff}$ ) but does not adequately capture the physics since the  $Z_{eff}$  dependence reduces the depth of the potential well to -55 V,

$$\Delta\varphi|_{eff} = \frac{T_e}{Z_{eff} \cdot T_e + T_i} \frac{m_{eff} \omega_\phi^2}{2e} (R^2 - R_0^{*2}) \quad (2)$$

flux-surface. This effect has already been observed in NSTX as is shown in Figure 2. The radiated power at the low-field-side (LFS,  $R=120$  cm) is higher than at its correspondent radii at the high field side (HFS,  $R \sim 80$  cm) possibly due to an asymmetric poloidal distribution of medium-Z impurities.

Due to the different effect of centrifugal forces




Fig. 4. a) Nested electron and ion temperatures and b) electron density offset from  $\psi(R,Z)$ .

The iterative solution for the electrostatic potential from the measured carbon density and central toroidal rotation using NSTX data is shown in Fig. 5-c) and can be calculated using arbitrary values of the core low-Z impurity strength factor which often exceeds unity ( $\alpha_{C,0} > 1$ ). The well's depth of the iteration solution is nearly 50% stronger than the ideal case and nearly twice as the *ad-hoc* approximation. A time-dependent solution (not shown here) shows also an in/out asymmetry evolution with strong  $\omega^2$  scaling as in eqns. (1) and (2). The depth of the potential well in the HFS can reach -110 to -280 V for core NSTX rotation velocities between 180 – 360 km/s but remains small if one compares with the core plasma potential (e.g. few kilovolts) or the energy of circulating and/or trapped fast ions (e.g. few tens of kilovolts).

The net-change of the plasma potential profile due to the presence of the rotation-induced electrostatic well is very small and of the order of 5-6% as is shown in Fig. 5-d). This small effect can be understood also analytically calculating the scaling of the changes of the radial electric field. The rotation-induced electrostatic potential can be written as in eqns. (3) and (4) with  $C$  being a constant that normalizes the numerical iteration solutions to the ideal approximation and  $L\omega\phi$  is the gradient scale length of the toroidal rotation frequency,

$$\Delta\varphi(r, \theta = \pi) \equiv \frac{CT_e}{T_e + T_i} \frac{m_i \omega_\phi^2}{2e} (R^2 - R_0^{*2}) \approx -2CR_0r \frac{T_e}{T_e + T_i} \frac{m_i}{e} \omega_\phi^2 \quad (3)$$

$$\Rightarrow \Delta E_r(r, \theta = \pi) \equiv -\nabla_r(\Delta\varphi(r, \theta = \pi)) = 2CR_0 \frac{T_e}{T_e + T_i} \frac{m_i}{e} \omega_\phi^2 (1 + 2r/L_{\omega\phi}) \quad (4)$$

The core electric field in NSTX is mainly determined by the  $V\phi B\theta$  term [7] which can be further simplified writing  $B\theta(r) = rB\phi/q(r)R$  and the toroidal field dependence as  $B\phi \sim B\phi_{,0}R_0/R$ , where  $B\phi_{,0}$  is the toroidal magnetic field at the magnetic axis,

$$E_r \approx V_\phi B_\theta \sim \frac{R_0 r^2}{q(r) R^2} \omega_\phi B_{\phi,0} \quad (5)$$

Using eqns. (4) and (5), the fractional change of radial electric field for  $T_e/(T_e + T_i) \sim 0.45$  is,

$$\Rightarrow \frac{\Delta E_r}{E_r}(r, \omega_\phi, \theta = \pi) \sim 0.9Cq(r) \left(\frac{R}{r}\right)^2 \frac{\omega_\phi(r)}{\Omega_{i,0}} (1 + 2r/L_{\omega\phi}) \quad (6)$$

where,  $\Omega_{i,0}$  is the core ion gyro-frequency. The ratio of the analytic expressions for the asymmetric radial electric field to the background radial field for  $C=1.55$ ,  $q(r)=1.4$ ,  $R \sim 80$  cm,  $r \sim 22.5$  cm,  $\omega\phi=2.15 \times 10^4$  Hz and  $\Omega_{i,0}=2.15 \times 10^7$  Hz, is of the order of 5% which is nearly the same as the one calculated experimentally and shown in Fig. 5-d).

#### IV. Impurity density and radiated power density profiles.

The computation of rotation-induced electrostatic potentials is also being used to study the associated two-dimensional distribution of impurity density and radiated power asymmetries




Fig. 5. a) Ideal, b) *ad-hoc* and c) iterative solutions of  $\Delta\varphi$  and its effect changing the mid-plane plasma potential.

and related change of the local electric field can be written as in eqns. (3) and (4) with  $C$  being a constant that normalizes the numerical iteration solutions to the ideal approximation and  $L\omega\phi$  is the gradient scale length of the toroidal rotation frequency,

in NSTX and NSTX-U. This calculation also finds 2D asymmetry factors (see Fig 6) for impurity profiles which are at the trace limit with very small changes to quasi-neutrality and  $Z_{\text{eff}}$ . Estimates of average charge state and radiated power density ( $P_{\text{rad}}$ ) can be made using coronal equilibrium. While the asymmetry in the core radiated power from low-Z ions (e.g. D, C, O, Ne) is relatively small, the core density and radiation from medium- to high-Z's will be strongly affected by centrifugal forces (see 2D  $P_{\text{rad}}$  profiles in Fig 7). At even higher toroidal rotation, the asymmetry is stronger concentrating the Fe, Mo and W radiated power strictly to the low-field-side.

In summary, the computation of rotation-induced electrostatic potentials is being used to study the associated two-dimensional distribution of impurity density asymmetries in NSTX. This calculation relies on flux-surface quantities like  $T_e$ ,  $T_i$  and  $\omega_\phi$ . The iterative process finds the 2D density profiles and the electrostatic potentials ( $\Delta\phi$ ) self-consistently assuming poloidal variation due to centrifugal forces. The depth of the potential well can reach -110 to -280 V for core NSTX rotation between 180 – 360 km/s but remains very small if one compares with the core plasma potential or the energy of fast ions. The net-change of the plasma potential and radial electric field is of the order of just 5-6% and in accordance with a simple theoretical calculation. This computation is being used to increase our understanding of asymmetries and the reduction of Z-peaking, to examine the effect of electrostatic potentials possibly changing the heat and particle transport, radiation asymmetries before tearing mode onsets, as well as to aid the design of new diagnostics for NSTX-U (e.g. ME-SXR [9], XICS, Bolometers, etc). This work is supported by the U.S. Department of Energy, Office of Fusion Energy Sciences under contract number DE-AC02-09CH11466.

## V. References.

[1] W. Houlberg, et al., PoP, **4**, 3320, (1997). [2] J. A. Wesson, NF, **37**, 577, (1997). [3] E. Belly and J. Candy, PPCF, **51**, 075018, (2009). [4] L. Delgado-Aparicio, et al., NF, **51**, 083047, (2011). [5] A. Mollen, et al., PoP, **19**, 052307, (2012). [6] T. Odstrcil, et al., PPCF, **60**, 014003, (2018). [7] R. Bell, PoP, **17**, 082507, (2010). [8] L. Delgado-Aparicio, to be submitted to NF, (2018). [9] L. Delgado-Aparicio, RSI, **87**, 11E204, (2016).




Fig. 7. Core impurity radiation from medium- to high-Z's (e.g. Argon, Iron, Molybdenum and Tungsten) will be strongly affected by centrifugal forces.




Fig. 6. Mass-dependence of asymmetry due to centrifugal forces. The depth of the potential well can reach -110 to -280 V for core NSTX rotation between 180 – 360 km/s but remains very small if one compares with the core plasma potential or the energy of fast ions.