

# Numerical simulation of thermal quench triggered by density source in HL-2A

S. L. Hu<sup>1</sup>, D. Hu<sup>2</sup>, J. Q. Li<sup>1</sup>, G. Z. Hao<sup>1</sup> & JOREK Team<sup>†</sup>

<sup>1</sup>*Southwestern Institute of Physics, PO Box 432, Chengdu 610041, China*

<sup>2</sup>*School of Physics, Beihang University No. 37 Xueyuan Road, Haidian District, Beijing 100083, China*

<sup>†</sup>*See the author list of Ref [5] for a list of current team members*

## Introduction

The mainstream methods of disruption mitigation system (DMS) include Shattered Pellet Injection (SPI) [1] or Massive Gas Injection (MGI) [2]. The injection penetration and density mixing are important considerations in achieving high mitigation efficiency, and they are strongly affected by MHD activities during injection. To clarify the effect of MHD-modes-induced-transport on the injection penetration, we simulate the thermal quench (TQ) during the pre-disruption phase triggered by pure deuterium ( $D_2$ ) injection at different fixed deposition location employing three-dimensional (3D) non-linear reduced MHD code JOREK [5].

## Physical model and simulation setting

Complete physics models implemented and numerical methods applied for solving the model equations in JOREK have been well documented [2-3]. We here only emphasize the essentials parameters relevant to our concerns. We refer to the equilibrium configuration of a typical Ohmic limiter discharge on HL-2A tokamak with  $B_t = 1.4$  T,  $I_p = 160$  KA. Central values approximately are  $n_0 = 1.38 \times 10^{19} \text{ m}^{-3}$ ,  $T_0 = 0.8$  kev. The resistivity and parallel thermal conductivity depend on temperature:  $\eta = \eta_0 (T_0/T)^{3/2}$  and  $\kappa_{\parallel} = \kappa_{\parallel 0} (T_0/T)^{5/2}$ , where  $\eta_0 = 5.3 \times 10^{-7} \Omega \text{m}$  is the Spitzer resistivity and  $\kappa_{\parallel 0} = 3.4 \times 10^{29} \text{ m}^{-1} \text{s}^{-1}$  is the Spitzer-Härm value in our simulation. Note that  $\eta_0$  and  $\kappa_{\parallel 0}$  closely approach to the experimentally measured values  $\eta_0 = 3.5 \times 10^{-7} \Omega \text{m}$ ,  $\kappa_{\parallel 0} = 2.0 \times 10^{29} \text{ m}^{-1} \text{s}^{-1}$ . Furthermore, we assume that constant  $10^{24} D$  atoms per second are injected.

Our simulations include all toroidal harmonics from  $n = 0$  to  $n = 5$  or  $n=0$  to  $n=10$ . To investigate the dependence of disruption mitigation on the location of D molecule disposition (LoD) with  $R = R_D$ , simulations are carried out with different penetration depth of D density




Fig 1. Flux surfaces before injecting deuterium, dotted lines represent rational surface.

source. Five typical cases are respectively  $R_D = R_j = 1.87, 1.91, 1.94, 1.97$  and  $2.02$  (m) at major radial direction, as shown in Fig.1 with  $j = (a, b, c, d, e)$  respectively. For initial equilibrium, the major radial position of the  $q = 1, 2, 3$  surfaces are respectively  $1.75, 1.92$  and  $2.0$  (m).

### Simulations on the TQ processes

After a long time steady state, active MHD fluctuation dominated by  $n=1$  mode grows up after the D density source is imposed in simulations. As the magnetic fluctuation increases, magnetic islands are created and grow. Afterwards, magnetic flux surfaces are destroyed and become stochastic. The core electron temperature rapidly decreases, turning into a TQ phase. The slight plasma current drop and the ensuing current spike is observed as the MHD perturbations develop. These typical processes can be shown in Fig.2, in which the results for the five cases mentioned above are exhibited for comparison. Interestingly, two types of evidently different disruption behaviors are observed, which depend on whether the D density source is deposited outside or inside the  $q=2$  magnetic surface.

The plasma current  $I_p$  and the fluctuating  $n=1$  magnetic energy reveals a remarkable spike in case  $R_c$ , which is one of characteristic features of plasma disruptions. However, the spike is much gentle in case  $R_a$ . This observation indicates that the  $q=2$  surface may be of crucial importance in triggering a major disruption by externally injected D density source.

We sample simulations with case  $R_c$  and case  $R_a$  as the targets for comparison and analysis. *The pre-TQ phase:*  $T_e$  at magnetic axis keep almost unchanged while the outer region are cooled down rapidly in both simulations. Significant increase of plasma density around the  $q=2$  surface on the LFS is observed in both cases, as  $R_c$  and  $R_a$  are located just outside and inside of said surface respectively. The island O-point coincides with the D deposition region, which is similar to the observation in [2, 4]. In this phase, it can be seen that the  $m/n = 2/1$  modes are excited around  $q=2$  surface in both simulations, but it is much stronger in case  $R_c$  than case  $R_a$ . Furthermore, in case  $R_c$ , the  $3/1$  magnetic islands also appear and overlap with the  $2/1$  magnetic




Figure 2. (a) Time evolution of the magnetic energy contained in the  $n=1$  component for five simulations (b) Time evolution of the  $I_p$  for the corresponding case. The dotted lines indicate respective the current spike time.

island. However, case  $R_a$  displays strong excitation of the 1/1 magnetic island.

*The TQ phase:* The  $T_e$  at the magnetic axis dramatically decreases, and plasma density further increases. In this stage, the MHD activity is relatively dominated by the  $m/n=2/1$  component in case  $R_c$  while the  $m/n=1/1$  mode dominates in case  $R_a$ . The TQ may result from successively nonlinear mode couplings, which are evidenced by local flattening of the toroidal current profile. Here, the local flattening projects the linear and/or nonlinear excitation of resonant mode. The 1/1, 2/1 and 3/1 mode are produced by MHD activity, then 2/1 mode is inward coupled with the 1/1 mode and outward coupled with the 3/1 mode. Afterwards, region near 3/2, 5/2 and 4/3 surfaces are flattened in case  $R_c$  while those around 3/2, 5/2 and 7/2 surfaces are flattened in case  $R_a$ . Therefore, the magnetic field tends to be stochastic in the region outside  $q=2$  surface and quickly expands inwards in case  $R_c$  while the magnetic field stochasticization occurs first in the region inside the  $q=2$  surface and expands outwards in case  $R_a$ . The TQ phase is completed at last and this stage is characterized by completely fallen-down of  $T_e$  and fully stochasticized magnetic field.

### Particle transport during pre-disruption phase

Most importantly, we found that increases of core density significantly depends on LoD and there exists a threshold, as shown in fig. 4 (a) by the average density profiles for five cases. It can be clearly seen that the core density in case  $R_a$  is much greater than other cases. The threshold is within the  $q=2/1$  surface. When




Figure 3. Toroidal current density  $j_{\phi,n=0}$  profiles at the midplane (low field side). (a) for case  $R_c$ , (b) for case  $R_a$ .




Figure 4. The average density profile, (a) five simulations in this text. The cyan short line is the  $q=1$  surface when the 1/1 mode grows up for  $R_a$ . (b) The initial  $q$  profile shifted down, two simulations:  $R_d = 1.935$  and  $1.975$  m.

the LoD crosses this threshold, the core density rises rapidly. To verify the effect of the  $q=1/1$  surface, we change initial equilibrium so that the  $q=1/1$  surface is shifted outward. Two cases are calculated, which corresponds to similar relative  $q$  profile position of case  $R_a$  and case  $R_c$ .

Results also show that increases of core density depend on LoD (seen from fig.4 (b)). The net radial particle fluxes are calculated in figure 5. It seems that increases of core density are due to  $\mathbf{E} \times \mathbf{B}$  convection produced by the 1/1 mode. We also compare the average particle fluxes produced by diffusion and convection inside the  $q=1/1$  surface. The result shows that  $\mathbf{E} \times \mathbf{B}$  convection is dominant as seen in table 1.

Table 1. The average particle flux inside minor radius

Position: (0.05-0.15) m unit:  $\times 10^{22} \text{m}^2/\text{s}$

	$Q_{n,DM}$	$Q_{n,(E \times B)r}$	$Q_{n,VparBr}$
$R_D=R_c$	0.15	-65.58	1.43
$R_D=R_a$	0.24	-271.88	3.29

## Conclusion

Some common features of the TQ process in disruption triggered by externally injected D density source have been observed by adjusting the LoD, including the MHD responses dominated by the  $n=1$  mode instabilities, a spike of plasma current and the magnetic field stochasticity first in the region of the  $m/n=2/1$  magnetic islands. Most importantly, it is shown that the TQ may result from successively nonlinear mode couplings. Comparisons between case  $R_c$  and case  $R_a$  reveal some evident differences depending on the LoD outside or inside the  $q=2$  surface. On one hand, the duration of the TQ phase is much shorter in case  $R_c$  than in case  $R_a$ , implying that a stronger major disruption could occur in case  $R_c$ . The magnetic fields are stochasticized first in the outer region of the  $q=2$  surface in case  $R_c$ , but in the inner region in case  $R_a$ . On the other hand, the core plasma density is visibly increased in case  $R_a$ . It seems that increases of core density are due to  $\mathbf{E} \times \mathbf{B}$  convection produced by the 1/1 mode.

- [1] D. Hu, E. Nardon et al., 45th EPS Conference on Plasma Physics, P4.1043.
- [2] E. Nardon, A. Fil et al., Plasma Phys. Control. Fusion 59, 014006(2017).
- [3] A. Fil, E. Nardon et al., Phys. Plasmas 22, 062509 (2015).
- [4] V. A. Izzo, Nucl. Fusion 46, 541–547(2006).
- [5] Matthias Hoelzl, Guido Huijsmans, et al., Nucl. Fusion 61, 065001 (2021).




Fig5. The particle flux before magnetic energy releasing, blue color is case  $R_c$  and red color is case  $R_a$ . The velocity expression is  $\mathbf{V} = -R^2 \nabla u \times \nabla \phi + v \parallel \mathbf{B}$ , The real line is produced by the  $r$  component of first term, and the dotted lines come from second term.